

Consultation Response Form

SCENIHR preliminary report on "The safety of dental amalgam and alternative dental restoration materials for patients and users"

Comment by

Prof. Vera Stejskal
Associated Professor of Immunology
University of Stockholm, Sweden and
Department of Immunology and Microbiology
1st Medical Faculty, Charles University, Prague, Czech Republic

Co-signed by

Prof. Marie Grosman
Prof. Agrégée de Biologie,
Association Non au mercure dentaire, FRANCE

Prof. André Picot
Directeur honoraire de Recherche CNRS
Président de l'Association de Toxicologie - Chimie ATC, Paris, FRANCE

Question 1: Is there scientific evidence that supports a link between amalgam and allergic reactions, neurological disorders or other health disorders?

Do you agree with the response given?

Disagree.

1. Unsatisfactory conclusion from the scientific point of view

2. Relevant information missing from the analysis of the situation

In addition to toxic effects, mercury induces local and systemic allergic and autoimmune reactions. Many metals, including mercury, function as haptens and induce cellular type hypersensitivity. This type of allergy is mediated by white blood cells (T-cells).

Inorganic mercury, thimerosal and nickel are the most frequent allergens in children as shown by skin patch test. In 1094 children with skin disease, 10% reacted to thimerosal (ethyl mercury salt) and 6 % to mercury (1). In 96 Spanish children, skin test reactivity to thimerosal was 21% and to mercury 19%. Body burden of mercury is associated with atopic eczema and total IgE antibodies in German children (3).

Below is a selection from the many articles indicating a causal relationship between mercury-induced sensitization and autoimmune diseases (4). The majority of patients improved following the removal of amalgam and other sensitizing dental restorations such as gold. The mechanisms behind of metal-induced effects in multiple sclerosis, rheumatoid arthritis and amyotrophic lateral sclerosis has been published by Stejskal and Stejskal (5).

Pelcova (6) reported skin exposure to mercury-containing creams which induced neuropsychological problems and glomerulonephritis in patients with juvenile diabetes. After chelation of mercury, the symptoms disappeared confirming a causal relationship.

Prochazkova (7) studied the impact of amalgam replacement on the health of patients with autoimmune diseases (multiple sclerosis, rheumatoid arthritis, psoriasis) who showed increased mercury-specific proliferation *in vitro*. Patients with only amalgam in the oral cavity were included in the study. Amalgam was replaced by composites and/or ceramics. Twenty out of 35 patients studied (71%) showed health improvement half a year later. Thus,

amalgam replacement might be beneficial in autoimmune patients with hypersensitivity to mercury.

Cellular hypersensitivity and autoantibodies to thyroid antigens were studied in 39 patients with autoimmune thyroiditis (8). Patients were divided into two groups, those with positive mercury-specific response *in vitro* and those with no stimulation with mercury *in vitro*.

Amalgam fillings were replaced in 15 patients with hypersensitivity to mercury and left in place in the remaining 12 patients (control group). Anti-thyroid peroxidase and anti-thyroglobulin antibodies were also measured. Only patients with mercury hypersensitivity who replaced their amalgam showed a significant decrease of autoantibodies compared to levels prior treatment. Thus, removal of amalgam in patients with mercury hypersensitivity might improve treatment of autoimmune thyroiditis. These results confirm the previous data (9-11). To our knowledge this is the first time when a specific biomarker of mercury susceptibility was used to select patients for amalgam replacement. Any risk factor may be diluted if evaluated in a heterogeneous population. As suggested by Weiss (12), studies of phenotypic markers may be suitable for elucidation of causal pathways, and identification of specific risk factors. The limited power of epidemiological studies to detect minor susceptible populations such as those susceptible to mercury has been discussed by Wallach (13) and Barregård (14).

Patch test and LTT-MELISA[®] were used for the diagnosis of metal allergy in 15 patients who suffered from clinical metal sensitivity and allergic and autoimmune diseases (15). The concordance of the two tests was good but the *in vitro* test was more sensitive. The removal of allergy-inducing dental restorations (amalgam and gold) resulted in long-term health improvement (follow up to 15 years). The improvement related to the decrease of metal-specific lymphocyte responses *in vitro*. Thus, in susceptible patients, metal ions might activate T-cells and start the inflammatory cascade. Replacement of inflammation-inducing materials results in decreased systemic inflammation and improved health.

References

1. Seidenari S, Giusti F, Pepe P, Mantovani L. Contact sensitization in 1094 children undergoing patch testing over a 7-year period. *Pediatr Dermatol* 2005;22:1-5
2. Vozmediano JMF, Hita A. Allergic contact dermatitis in children. *J European Academy Dermatol venerol* 2005;19:42-46
3. Weldinger S, Krämer U, Dunemann L, Möhrenscklager M, Ring J et al. *J Allergy Clin Immunol* 2004;114:457-459
4. Cooper G, Germolec D, Heindel, Selgrade. Linking environmental agents and autoimmune diseases. *Environ health Persp* 1999;107 (Suppl 5): 659-660.
5. Stejskal J, Stejskal V. The role of metals in autoimmunity and the link to neuroendocrinology. *Neuro Endocrinol Lett* 1999;20:351-364
6. Pelcova D, Lukas E, Urban P, Preiss J, Rysava R et al. Mercury intoxication from skin ointment containing mercuric ammonium chloride. *Int Arch Occup Environ Health* 2002;75(Suppl): S54-S59
7. Prochazkova J, Sterzl I, Kucerova H, Bartova J, Stejskal V. The beneficial effect of amalgam replacement on health in patients with autoimmunity. *Neuro Endocrinol Lett*. 2004;25:211-8.
8. Sterzl I, Prochazkova J, Hrda P, Matucha P, Bartova J et al. Removal of dental amalgam decreases anti-TPO and anti-Tg autoantibodies in patients with autoimmune thyroiditis. *Neuro Endocrinol Lett* 2006;27:25-30

9. Sterzl I, Prochazkova J, Hrda P, Bartova J, Matucha P, Stejskal VDM. Mercury and nickel allergy: risk factors in fatigue and autoimmunity. *Neuro Endocrinol Lett*. 1999;20:221-228.
10. Lindkvist B, Mörnstad H. Effects of removing amalgam fillings from patients with diseases affecting the immune system. *Med Sci Res* 1996;34:355-356
11. Anneroth G, Ericson T, Johansson I, Mörnstad H, Ryberg M et al. Comprehensive medical examination of a group of patients with alleged adverse effects from dental amalgam. *Acta Odont Scand* 1992;50:101-111
12. Weiss NS, Liff JM. Accounting for the multicausal nature in disease in the design and analysis of epidemiological studies. *Am Epidemiol* 1983;117:14-18.
13. Wallach H, Nauman J, Mutter J, Daschner F. No difference between self-reportedly amalgam sensitivite and non-sensitivites? Listen carefully to the data. *Int J Hyg Environ Health* 2003; 206:139-141
14. Barregård L, Eneström S, Ljunghusen O, Wieslander J, Hultman P. A study of autoantibodies and circulating immune complexes in mercury-exposed chloralkali workers. *Int Arch Occup Environ Health* 1997; 70:101-6
15. Stejskal V, Hudecek R, Stejskal J, Sterzl I. Diagnosis and treatment of metal-induced side-effects *Neuroendo Lett* 2006;27(Suppl 1):7-16

More references on the subject not mentioned in the text due to limited space:

- Balasz T.** Immunogenetically controlled autoimmune reactions induced by mercury, gold and D-penicillamine in laboratory animals: a review from the vantage point of premarketing safety studies. *Toxicol and Industrial health* 1987;3:331-336
- Bangsi D** et al. Dental amalgam and multiple sclerosis: a case-control study in Montreal, Canada. *Int J Epidemiol* 1998;27:667-71
- Bigazzi PE** Autoimmunity and heavy metals. *Lupus* 1994;3:449-453
- Bigazzi PE.** Metals and kidney autoimmunity. *Environ Health Persp* 1999;107 (Suppl5):753-765
- Campbell A** et al. Mechanisms by which metals promote events connected to neurodegenerative diseases. *Brain Res* 2001;55:125-132
- Cooper GS** et al. Occupational risk factors for the development of systemic lupus erythematosus. *J Rheumatol* 2004;31:1928-33
- Charpentier B** et al. Fonctions lymphocytaires T au cours d'une glomerulonephrite extramembraneuse induite par une intoxication chronique au mercure. *Nephrologie* 1981;2:153-157
- Druet P.** Metal-induced autoimmunity. *Human and Expt Toxicol* 1995;14:120-121
- El-Fawal HA** et al. Neuroimmunotoxicology: humoral assessment of neurotoxicity and autoimmune mechanisms. *Environ Health Persp* 1999;107:767-75
- El Safty IA** et al. Nephrotic effects of mercury exposure and smoking among Egyptian workers in fluorescent lamp factory. *Arch Med Res* 2003;34:50-55
- Hock C** et al. Increased blood mercury levels in patients with Alzheimer's disease. *J of Neural Trans* 1998;105:59-68
- KazantisG.** Mercury exposure and early effects: an overview. *Med Lav* 2002; 93:139-46
- Kosuda LL** et al. Effects of HgCl₂ on the expression of autoimmune responses and disease in diabetes-prone (DP) BB rats. *Autoimmunity* 1997;26:173-187
- Lindh U** et al. Removal of dental amalgam and other metal alloys supported by antioxidant therapy alleviated symptoms and improves quality of life in patients with amalgam-associated ill health. *Neuroendo Lett* 2002;23:459-482
- Mach E** et al. Umweltgifte und multiple Sklerose. *Der Allgemeinartz* 1996;20:2216-2219
- Queiroz R** et al: Immunoglobulin levels in workers exposed to inorganic mercury. *Pharmacol Toxicol* 1994;74:72-75

- Shenker BJ** et al. Immunotoxic effects of mercuric compounds on human lymphocytes and monocytes III. Alterations in B-cell function and viability. *Immunopharmacol Immunotoxicol* 1993;15:87-112
- Silbergeld EK** et al. Mercury and autoimmunity: implications for occupational and environmental health. *Toxicol Appl Pharmacol* 2005;207:282-92
- Siblerud RL.** A comparison of mental health of multiple sclerosis patients with silver/mercury fillings and those with fillings removed. *Psychol Rep* 1992;70:1139-51
- Silva IA** et al. Mercury exposure malaria and serum antinuclear/antinucleolar antibodies in Amazon populations in Brazil: a cross-sectional study. *Environ Health* 2004;3:11
- Tubbs RR** et al. Membranous glomerulonephritis associated with industrial mercury exposure. Study of pathogenic mechanisms. *Am J Clin pathol* 1982;77:409-13
- Tchounwou PB,** et al. Environmental exposure to mercury and its toxicopathologic implications for public health. *Environ Toxicol* 2003;18:149-75
- Valentin-Thon E** et al. Validity of MELISA[®] for metal sensitivity testing. *Neuroendo Lett* 2003; 24:57-64
- Via Ch** et al. Low-dose exposure to inorganic mercury accelerated disease and mortality in acquired murine lupus. *Environ Health Perspectives* 2003;111:1273-1277

Question 2: In view of the above, is the use of dental amalgam safe for patients and users, i.e. dental health professionals? Are certain populations particularly at risk, e.g. pregnant women or children?

Disagree

- 1. Unsatisfactory conclusion from the scientific point of view**
- 2. Relevant information missing from the analysis of the situation**

Heavy metals including mercury are biologically active substances and may in susceptible subjects affect many organs and cause health disturbances. Heavy metals are known to induce so called cellular type hypersensitivity (delayed type or Type 4 reaction) but humoral antibodies might be affected as well. Metal-induced reactions are influenced by genetic background in experimental animals and associated with certain HLA antigens in man (1). Patients with allergic and autoimmune diseases such as multiple sclerosis, collagenous diseases, and psoriasis might be particularly vulnerable (2).

Mercury has been documented to be a reproductive and developmental toxin in humans. The effects include male infertility (3,4), lowered sperm counts, defective sperm cells, menstrual disturbances, infertility, spontaneous abortions and birth effects. Mercury causes learning disabilities and impairment and reduction in IQ. Regarding children, amalgam-treated children exhibited significantly higher microalbuminuria compared to children without amalgam (5).

The assumption that mercury released from amalgam only rarely induces allergy is wrong and is based on the observations of oral mucosal problems which are less frequent due to lower sensitivity of oral mucosa. In the oral cavity, a high concentration of metal ions may be toxic to immuno-competent cells and act as a local immunosuppressant. Oral mucosa contains only a low number of dendritic cells, and mucosal changes adjacent to dental metal fillings are infrequent (6). Nielsen and Klaschka (7) have shown that a 5-12 times higher concentration of the allergen has to be applied on the oral mucosa than on the skin to elicit microscopic reactions.

The authors of the SCENIHR report claim that it is not necessary to remove clinically satisfactory amalgam restorations on the grounds of patient safety, with the exception of those patients which have a positive patch test and local alterations of the oral mucosa or **systemic allergic reactions**. We agree with that. As mentioned in answer to Question 1, vulnerable groups includes children and adults with diseases of immune origin such as contact dermatitis and autoimmunity. By definition, in those patients, the immune system reacts aberrantly and might recognize mercury as a hapten and trigger allergic and autoimmune disease (8, 9). Therefore, amalgam has to be removed.

References

1. Prochazkova J, Bartova J, Ivaskova E, Kupka L, Sterzl I, HLA-association in patients with intolerance to mercury and other meals. *Disease Markers* 2001;16:135-138
2. Bartova J, Prochazkova J, Kratka Z, Benetkova K, Venclikova Z et al. Dental amalgam as a risk factor in autoimmune diseases. *Neuroendo Lett* 2003; 24:65-67
3. Choy CM Lam CV, Cheung LT, Briton-Jones CM et al. Infertility, blood mercury concentrations and dietary seafood consumption. *BJOG* 2002; 109:1121-1125
4. Hanf V, Forstman A, Costea JF, Schierferstein G, Fisher I et al. Mercury in urine and ejaculate in husbands of barren couples, *Toxicol lett* 1996;1-3:227-231
5. Barregard L, Trachtenberg F, Mc Kinlay S. Effects of Dental Amalgam in Children: The New England Children's Amalgam Trial *Environ Health Perspect* 2008;116:394–399
6. Magnusson B, Bergman M, Bergman B, Sörenmark R. Nickel allergy and nickel-containing dental alloys. *Scand J Dent Res.* 1982;90:163-65
7. Nielsen C, Klaschka F. Test studies on mouth mucosa in allergic eczema. *Dtsch Zahn Mund Kieferheilkd Zentralbl Gesamte* 1971;57:201-18
8. Stejskal V, Hudecek R, Stejskal J Sterzl I. Diagnosis and treatment of metal-induced side-effects. *Neuroendo Lett* 2006;27(Suppl 1):7-16
9. Stejskal J, Stejskal V. The role of metals in autoimmunity and the link to neuroendocrinology. *Neuro Endocrinol Lett* 1999;20:351-364

Question 3: Is there scientific evidence that supports a link between alternative materials and allergic reactions, neurological disorders or other health disorders?

YOUR COMMENTS

Do you agree with the response given?

Mostly agree, regarding non-metallic materials..

Question 4: In view of the above, is the use of alternative dental restoration treatment safe for patients and dental health professionals? Are certain populations particularly at risk, e.g. pregnant women or children?

YOUR COMMENTS

Do you agree with the response given?

Yes (non metallic restorations)

No (gold, nickel and titanium alloys).

1. Unsatisfactory conclusion from the scientific point of view

2. Relevant information missing from the analysis of the situation

Regarding metallic restorations for example gold alloys, nickel alloys and titanium alloys they contain transitional metals which may in susceptible subject trigger allergy and autoimmunity (1-12). Gold is now the second most common sensitizer in man after nickel. Palladium, as well as titanium, is a transition metal with the capacity to bind to proteins and cause sensitization (13-15). Regarding non metallic materials such as composites and ceramics, we agree with the response given by authors.

References:

1. Stejskal V, Hudecek R, Stejskal J Sterzl I. Diagnosis and treatment of metal-induced side-effects. *Neuroendo Lett* 2006;27(Suppl 1):7-16
2. Magnusson B, Bergman M, Bergman B, Södermark R. Nickel allergy and nickel-containing dental alloys. *Scand J Dent Res* 1992;90:163-167
3. Samitz MH, Katz SA. Nickel dermatitis hazards from prostheses. *British J Dermatol* 1975;92:287-290
4. Bruce GJ. Nickel hypersensitivity-related periodontitis. *Compendium* 1995;16:178-184
5. Marcusson JA Lindh G, Evengård B. Chronic fatigue syndrome and nickel allergy. *Cont Derm* 1999;40:269-272
6. Björkner B, Bruze M, Möller H. High frequency of contact allergy to gold thiosulfate. An indication of gold allergy? *Contact Derm* 1994;30:144-151
7. Marcusson JA. Contact allergies to nickel sulfate, gold sodium thiosulfate and palladium chloride in patients claiming side-effects from dental alloy components. *Contact Derm* 1996;34:320-323
8. Schlumpf, Meyer J, Ulrich, Friede RL. Neurologic complications induced by gold treatment. *Arthritis Rheumat* 1983;26:825-831
9. Agrup G. Sensitization induced by patch testing. *Br J Derm* 1968;80:631-634
10. Räsänen L, Tuomi ML. Diagnostic value of the lymphocyte proliferation test in nickel contact allergy and provocation in occupational coin dermatitis. *Contact Derm* 1992;27:250-2
11. Muris J, Feilzer A. Microanalysis of metals in dental restorations as part of diagnostic approach in metal allergies. *Neuroendo Lett*: 2006;27(Suppl1):49-52
12. Venclikova Z, Benada O, Bartova J, Joska L, Mrklas L et al. In vivo effects of dental casting alloys. *Neuroendo Lett*; 2006;27(Suppl1):61-68
13. Aberer W, Holub H, Strohal L, Slavicek L. Palladium in dental alloys: the dermatologist's responsibility to warn? *Contact Derm* 1998;28:163-168
14. Valentine-Thon, Muller K, Guzzi G, Kreisel S, Ohnsorge P et al. LTT-MELISA® is clinically relevant for detecting and monitoring metal sensitivity. *Neuroendo Lett*; 2006;27(Suppl1):17-24
15. Muller K, Valentine-Thon E. Hypersensitivity to titanium: clinical and laboratory evidence. *Neuroend Lett*; 2006;27(Suppl1):31-35

Question 5: In view of the specific properties of dental amalgam and alternatives when used for dental restorative treatment, is dental health equally ensured by dental amalgam and alternatives?

Do you agree with the response given?

Disagree

1 Unsatisfactory conclusion from the scientific point of view

2. Relevant information missing from the analysis of the situation

The authors of SCENIHR report claim that they see no advantages to carrying out further research on any aspects of the safety of dental amalgam restorations.

We disagree. More research is necessary, especially prospective longitudinal studies in susceptible subjects. Since it is not ethical to insert amalgams to children with already compromised immune systems (those with allergies and autoimmunity), longitudinal studies are necessary when careful replacement of amalgam with ceramic and composite materials

will be performed and the health outcome monitored. Such treatment can be done in addition to standard therapeutic treatment for the disease in question and compared to the treatment without the replacement of amalgam (and other sensitizing metals in question). More research is also necessary to identify the biomarkers of susceptibility at the immunological and biochemical level. For example, biomarkers of harmful effects of metals and other environmental pollutants include detoxification enzymes, such as apolipoprotein E, where the substitution of cysteine with arginine – an amino acid lacking SH-groups – predisposes for increased risk for Alzheimer's disease (1) and increases vulnerability to chronic mercury toxicity (2). Other detoxification enzymes of importance are glutathione S transferase T1 (GSTT1) and glutathione S transferase M1 (GSTM1). As shown by Westphal's group (3), homozygous deletion of GSTT1 and combined deletion of GSTT1-/GSTM1- was markedly more frequent in patients sensitized by thimerosal, than in healthy controls. Regarding metal susceptibility, measurement of beryllium specific memory cells in the blood of exposed workers is currently the golden standard for detection of beryllium susceptibility (4-6). We postulate that a similar approach should be used for screening of patients at risk for side-effects of dental material.

In conclusion, susceptible populations at risk due to mercury and other metals are children and adults with allergic and autoimmune diseases. Children with autistic and behavioral disorders belong with all probability to the susceptible group as well. Until now, epidemiological studies either excluded these groups (7) or had limited power to detect those risks (8,9).

In the future, the best way to study the possible role of metals in the pathogenesis of diseases seems to be:

- 1) Selection of susceptible patients on the basis of phenotype and genotype from the heterogeneous cohort**
- 2) Therapy based on the elimination of the exposure to putative allergen(s)**
- 3) Long-term follow-up of patient's health combined with monitoring of improvement in relevant laboratory markers.**

References

1. Godfrey ME, Wojcik DP, Krone CA. Apolipoprotein E genotyping as a potential biomarker for mercury neurotoxicity. *J Alzheimers Dis.* 2003;5:189-95.
2. Wojcik DP, Godfrey ME, Christie D, Haley BE. Mercury toxicity presenting as chronic fatigue, memory impairment and depression: diagnosis, treatment, susceptibility, and outcomes in a New Zealand general practice setting (1994-2006). *Neuro Endocrinol Lett.* 2006;27:415-23
3. Westphal GA, Schnuch A, Schulz TG, Reich K, Aberer W, Brasch J, et al. Homozygous gene deletions of the glutathione S-transferases M1 and T1 are associated with thimerosal sensitization. *Int Arch Occup Environ Health.* 2000;73:384-8)
4. Kreiss K, Newman LS, Mroz M, Campbell PA. Screening blood test identifies subclinical beryllium disease. *J Occup Med.* 1989;31:603-8,
5. Kreiss K, Miller F, Newman LS, Ojo-Amaze E, Rossman MD, Saltini C. Chronic beryllium disease – from the workplace to cellular immunology, molecular immunogenetics, and back. *Clin Immunol Immunopathol.* 1994;71:123-9
6. Newman LS. Significance of the blood beryllium lymphocyte proliferation test. *Environ Health Perspect.* 1996;104(Suppl 5):953-6
7. Bellinger DC, Trachtenberg F, Barregard L, Tavares M, Cernichiari E et al. Neuropsychological and renal effects of dental amalgam in children. A randomized clinical trial *JAMA* 2006;295:1775-1783

8. Wallach H, Nauman J, Mutter J, Daschner F. No difference between self-reportedly amalgam sensitivitie and non-sensitivites? Listen carefully to the data. *Int J Hyg Environ Health* 2003;206:139-141
9. Barregård L, Eneström S, Ljunghusen O, Wieslander J, Hultman P. A study of autoantibodies and circulating immune complexes in mercury-exposed chloralkali workers. *Int Arch Occup Environ Health* 1997;70:101-6